



Environmental hazard classification and labelling

Jonas Falck
Swedish Chemicals Agency


ITP 22 September 2014

Contents

- Background and scope
- The classification scheme
- Criteria for env. hazard classification – substances
- Degradation and Bioaccumulation assessment for classification purposes
- Exercise – substance classification
- Criteria for env. hazard classification – mixtures
- Exercise – mixture classification
(principle use of the Summation method)

Ecotoxicological concept

- Ecology
- Toxicology
- Environmental Chemistry



Ecotoxicology

- Concerned with **adverse effects** of chemical and physical agents on living organisms, especially on populations and communities **within defined ecosystems**.

Environmental hazard classification – Define effects on **ecosystems** rather than on individuals within a species or population. E.g. Hazardous to the aquatic environment

- **Short-term (Acute)** and **long-term** adverse effects

Example of an acute (and obvious) effect



Cyanide in spillage water from a goldmine in Rumania, 2000, caused severe fish death. Also rivers in Serbia were affected.

Examples of observed long term toxic effects in the environment



05F0D-00160565-891 [RM] © www.visualphotos.com

- Industrial melanism of moths - 1850s Industrial revolution soot from coal burning



- Eggshell thinning in eagles and brown pelicans - 1950s DDT and organo-chlorines

➤ The classification scheme

- is principally concerned with the **aquatic environmental compartment** (which for most substances, the majority of data available addresses)

➤ This compartment is

- vulnerable
- receiving environment
- sensitive organisms



Scope

➤ The classification scheme covers both:

- short term effects
- long term effects

to both


- aquatic freshwaters
- marine ecosystems



What about the other compartments?

(E.g. the terrestrial compartment)

➤ Terrestrial test organisms



Not covered in a hazard classification scheme yet, but many substances hazardous to aquatic ecosystem would also be hazardous to terrestrial ecosystems.

Hazardous to the Ozone Layer

Substances

- if the available evidence concerning its properties and its predicted or observed environmental fate and behavior indicate that it may present a danger to the structure and/or the functioning of the stratospheric ozone layer.

Mixtures


- Concentration limit of 0.1%



WARNING

H420: Harms public health and the environment by destroying ozone in the upper atmosphere

What effects can be observed?



- Increased ecological relevance
- Increased difficulty to relate to a specific chemical
- Increased time from disturbance to effect

pragmatic choice:

- controlled conditions
- low natural variance,
- short time frame
- easy to observe
- cheap
- comparable between substances

For aquatic hazard classification, toxicity data is normally needed on three trophic levels



Fish



Crustacean





Alge/aquatic plants

- The taxa chosen from **three trophic levels** represent the “base-set” of toxicity test data; a minimum data-set for a fully valid description of toxicity as part of aquatic hazard.

Food chain with different trophic levels

Top-consumers




Secondary consumers

Primary consumers

Producers

Food web



A simplified food web for the Northwest Atlantic

Test methods for environmental toxicity and fate

- Test methods are **highly standardized**.
 - OECD test guidelines
 - EU test methods (Council regulation 440/2008)
 - ISO standards (CEN)
 - National: ASTM (USA), MITI (Japan), SIS (Sweden)
 - IOBC-guidelines and SETAC guidelines regarding arthropods

For aquatic hazard classification OECD Test Guidelines or equivalent, Ex.:

➤ **Physico-chemical properties:**

- 105 (Water solubility);
- 107 (n-octanol/water partition coefficient (Log K_{ow}))
- 111 (Hydrolysis as function of pH - Abiotic degradation)

➤ **Aquatic toxicity:**

- 201 (Algal Growth Inhibition);
- 202 Part 1&2, 211 (*Daphnia* sp. Acute Immob. & Reproduction);
- 203 (Fish, Acute Toxicity Test);
- 210 (Fish Early Life Stage)

➤ **Degradation:**

- 301A-F, 306, 310 (Ready biodegradability);
- 309 (Aquatic simulation test)

➤ **Bioaccumulation:**

- 305 (Bioconcentration factor in fish, BCF);

www.kemi.se

- Use of non testing methods -

➤ In absence of experimental data, valid non testing methods can be relied upon:

- Read across from similar chemicals
- Information from Chemical Structure - Structure-activity relationship (SAR)

Ex. provide predictions of acute toxicity by use of QSARs for:

- Non-electrolyte, non-electrophilic, and otherwise non-reactive organic substances.

e.g. hydrocarbons, alcohols, ketones and certain aliphatic chlorinated hydrocarbons and otherwise non-reactive substances

The classification and labelling schemes

Classification categories for Hazardous to the aquatic environment

See GHS,
Table 4.1.1

Hazard Class

Hazard Category

Hazardous to the aquatic environment

- Short-term (acute) hazard
- Long-term (chronic) hazard

Acute 1

Acute 2 *

Acute 3 *

Chronic 1

Chronic 2

Chronic 3

+ Chronic 4

NOTE!

Acute 1 to 3 + Chronic 1 to 3: The core classification system.

Chronic 4: 'Safety Net' classification when standard criteria are not met, but there is a concern.

Criteria: Not strictly defined, but one example: poorly soluble substances (< 1 mg/l) that are both

- not rapidly degradable and
- Bioaccumulative.

Short-term and Long-term hazard: are applied independently.

Classification categories for Hazardous to the aquatic environment

See GHS, Table 4.1.1

Hazard Class

Hazard Category

'cut offs'
≤ 1 mg/l

Hazardous to the aquatic environment

- Short-term (acute) hazard
- Long-term (chronic) hazard

Acute 1	Acute 2 *	Acute 3 *	
Chronic 1	Chronic 2	Chronic 3	+ Chronic 4


Relevant concentrations in the environment

Supply and use sector: ≤ 1 mg/l

Transport sector: ≤ 100 mg/l



Labelling elements

- Acute (short-term) aquatic hazard - Categories Acute 1 to 3

	<u>Acute 1</u>	<u>Acute 2</u>	<u>Acute 3</u>
Pictogram		No Pictogram	No Pictogram
Signal word	Warning	No word	No word
Hazard Statement	H400: Very toxic to aquatic life	H401: Toxic to aquatic life	H402: Harmful to aquatic life

Labelling elements

➤ Long-term aquatic hazard - Categories Chronic 1 to 3

	<u>Chronic 1</u>	<u>Chronic 2</u>	<u>Chronic 3</u>
Pictogram			No Pictogram
Signal word	Warning	No word	No word
Hazard Statement	H410: Very toxic to aquatic life with long-lasting effects	H411: Toxic to aquatic life with long-lasting effects	H412: Harmful to aquatic life with long-lasting effects

Safety net Chronic 4 - H413: May cause long lasting harmful effects to aquatic life.

Criteria for
environmental hazard
classification
-
substances

Acute toxicity determines the short-term hazard

In principle


- Intrinsic property to be injurious in short-term exposure
 - (hours to days)
- Generally expressed:
 - LC_{50} (50% lethal conc.) or EC_{50} (50% effect conc.),
e.g. immobilization of daphnids, or reduction in growth rate in algae

Chronic toxicity determines the long-term hazard

determined by

- Intrinsic property to be injurious during exposures which are determined in relation to the life-cycle of the organism
 - (days to weeks)
- Generally expressed in terms of:
 - NOEC, LOEC or EC_x (Normally EC_{10})
Sublethal endpoints e.g. Survival, growth and/or reproduction

Dose-response relationship



*No Observed
Effect Concentration*

*Concentration causing
effect on 50% of test
organisms*

Acute (short-term) aquatic hazard

Highest acute toxicity (lowest value) to
● Fish ● Crustacea or ● Aquatic plant

➤ Category

LC_{50} or EC_{50} (or IC_{50}) ≤ 1 mg/l

Acute 1

LC_{50} or EC_{50} (or IC_{50}) > 1 to ≤ 10 mg/l

Acute 2 *

LC_{50} or EC_{50} (or IC_{50}) > 10 to ≤ 100 mg/l

Acute 3 *


* Categories Acute 2 and 3 were mainly meant for transport of bulk-quantities and therefore normally not implemented for Supply & Use

Basic elements used for Long-term hazard

See GHS,
Table 4.1.1

Chronic toxicity data are often expensive to generate and therefore generally less available than acute toxicity data.

- For practical reasons a limited set of specific properties (basic elements) has been selected through which the hazard can be best described.



Criteria for Long-term hazard

the "surrogate system"

Adequate chronic toxicity data available		In absence of adequate chronic toxicity data
Non-rapidly degradable (NRD) substance	Rapidly degradable (RD) substances	ACUTE TOXICITY + NON-RAPIDLY DEGRADABLE and/or BIOACCUMULATIVE
Category: Chronic 1 NOEC or $EC_x \leq 0.1$	Category: Chronic 1 NOEC or $EC_x \leq 0.01$	
Category: Chronic 2 $0.1 < NOEC$ or $EC_x \leq 1$	Category: Chronic 2 $0.01 < NOEC$ or $EC_x \leq 0.1$	
	Category: Chronic 3 $0.1 < NOEC$ or $EC_x \leq 1$	

Regulatory acceptance based on relevant concentrations in the environment

Toxicity + degradation and/or bioaccumulation

A/C = 10 and 100

Long-term hazard in absence of adequate chronic toxicity data

**ACUTE TOXICITY TO FISH
CRUSTACEA OR ALGAE**

≤ 1 mg/l

+

**NON-RAPIDLY
DEGRADABLE**



CHRONIC 1

**ACUTE TOXICITY TO FISH
CRUSTACEA OR ALGAE**

> 1 to ≤ 10 mg/l

+

AND/OR



CHRONIC 2

**ACUTE TOXICITY TO FISH
CRUSTACEA OR ALGAE**

> 10 to ≤ 100 mg/l

+


**BIOACCUMULATIVE
(measured
or potential for
bioaccumulation)**




CHRONIC 3

See GHS, Figure 4.1.1


Adequate Chronic toxicity data



Classify according to the criteria given ... or ... depending on inf. On rapid degr.



Classify both:
a) according to the criteria given ... or ... depending on inf. On rapid degr.
and
b) For other trophic levels, the "surrogate system"
and classify according to the most stringent outcome



Classify according to the criteria given ... (the "surrogate system")

Criteria for Long-term hazard

Adequate chronic toxicity data available		In absence of adequate chronic toxicity data
Non-rapidly degradable substances	Rapidly degradable substances (RD)	
Category: Chronic 1 NOEC or $EC_x \leq 0.1$	Category: Chronic 1 NOEC or $EC_x \leq 0.01$	Category: Chronic 1-3 ACUTE TOXICITY + BIOACCUMULATIVE and/or LACK OF RAPID DEGRADATION
Category: Chronic 2 $0.1 < \text{NOEC or } EC_x \leq 1$	Category: Chronic 2 $0.01 < \text{NOEC or } EC_x \leq 0.1$	
	Category: Chronic 3 $0.1 < \text{NOEC or } EC_x \leq 1$	
Safety net classification Category: Chronic 4 When standard criteria are not met, but there is a concern. Not strictly defined criteria, but one example: <u>poorly soluble substances</u> ($< 1 \text{ mg/l}$) that are <u>not rapidly degradable</u> and are <u>bioaccumulative</u>		

M-factors must be set for highly toxic substances

See GHS,
Note 2 to Table 4.1.1

Hazard Class

Hazard Category

Hazardous to the aquatic environment

- Short-term (acute) hazard

Acute 1

Acute 2 *

Acute 3 *

- Long-term (chronic) hazard

Chronic 1

Chronic 2

Chronic 3

+ Chronic 4

- 'M-factor' means a multiplying factor. It is applied to substance as part of the substance classification as Categories Acute 1 and/or Chronic 1.
- It is used to derive by the summation method the classification of a mixture in which the substance is present.

Setting M-factors for highly toxic substances (Acute 1 and Chronic 1)

Acute toxicity	M factor
L(E)C ₅₀ value (mg/l)	
$0.1 < L(E)C_{50} \leq 1$	1
$0.01 < L(E)C_{50} \leq 0.1$	10
$0.001 < L(E)C_{50} \leq 0.01$	100
$0.0001 < L(E)C_{50} \leq 0.001$	1000
$0.00001 < L(E)C_{50} \leq 0.0001$	10000
(continue in factor 10 intervals)	

Setting M-factors for highly toxic substances (Acute 1 and Chronic 1)

Acute toxicity	M factor	Chronic toxicity	M factor	
L(E)C ₅₀ value (mg/l)		NOEC value (mg/l)	NRD ^a components	RD ^b components
$0.1 < L(E)C_{50} \leq 1$	1	$0.01 < NOEC \leq 0.1$	1	-
$0.01 < L(E)C_{50} \leq 0.1$	10	$0.001 < NOEC \leq 0.01$	10	1
$0.001 < L(E)C_{50} \leq 0.01$	100	$0.0001 < NOEC \leq 0.001$	100	10
$0.0001 < L(E)C_{50} \leq 0.001$	1000	$0.00001 < NOEC \leq 0.0001$	1000	100
$0.00001 < L(E)C_{50} \leq 0.0001$	10000	$0.000001 < NOEC \leq 0.00001$	10000	1000
(continue in factor 10 intervals)		(continue in factor 10 intervals)		

Degradation and Bioaccumulation assessment for classification purposes

Rapid degradation

- biotic or abiotic
 - degradation of organic substances; or
 - transformation of inorganic substances
- Either
 - full mineralisation or
 - primary degradation / transformation to non hazardous species ($t_{1/2} < 16$ days)



Rapid degradation - Decision scheme

GHS Annex 9
A9.4.4

A substance is considered to be **not** rapidly degradable **unless** at least one of the following is fulfilled:

- a) Ultimately degraded in biodegradation **screening test** ($\geq 60/70\%$ in 28days);
- b) Ultimately degraded in a surface water **simulation test** ($t_{1/2} < 16$ days);
- c) Primarily degraded (or transformed) to non hazardous species ($t_{1/2} < 16$ d)

When these **preferred data** types are **not available** rapid degradation may be demonstrated if one of the following criteria is justified:

- a) Ultimately degraded in an aquatic sediment or soil simulation test;
- b) If only BOD5 and COD available, then if $BOD5/COD \geq 0.5$;
- c) A weight of evidence approach based on read-across

If none of the above types of data are available then the substance is considered as **not** rapidly degradable.

Biodegradation Screening test vs. Simulation tests

Screening tests

- **Tests conducted in the laboratory with relatively high concentrations of test substance (2-100 mg/l).**
- All organic substances that degrade to a level higher than the pass level in a standard ready biodegradability test (OECD 301 A-F, 306 and 310 or similar test) should be considered rapidly degradable.
 - ≥ 70 %, 28-day test, based on dissolved organic carbon
 - ≥ 60 %, 28 day test, O₂-depletion or CO₂-generation

Biodegradation Screening test vs. Simulation tests

Simulation tests

- **Tests conducted in the laboratory, but simulating environmental conditions and employing natural samples as inoculum.**
- An environmental simulation test would normally be conducted according to one or more of the standard procedures of OECD Guidelines:
 - 307 (soil),
 - 308 (aquatic sediment), or
 - **309 (water)**

Biotic vs. abiotic degradation

Hydrolysis (abiotic degradation)

- Data on hydrolysis might be considered for classification purposes to measure the longest half-life $t_{1/2}$ determined within the pH range 4 - 9.
- E.g. OECD 111.





Degradation data not used for classification

- Inherent biodegradability (e.g. OECD 302)
- Sewage treatment plant (STP) simulation tests (e.g. OECD 303)
- Anaerobic degradation data
- Field investigations
- Monitoring data
- Photochemical degradation
- Volatilisation

Bioaccumulation

Def.: The net result of uptake, transformation and elimination of a substance in an organism

- Generally expressed in terms of:
 - Bioconcentration factor ($BCF \geq 500$),
(The ratio between the conc. in biota and the conc. in surrounding medium, pref. whole fish/water, and
 - in absence of BCF, the Octanol-water-partitioning coefficient ($\log K_{ow} \geq 4$)



ECHA guidance documents

➤ **Introductory Guidance on the CLP Regulation**

- Basic guidance for inexperienced classifiers and managers;


Explains the system (roles and obligations)
and why we have it.



➤ **Guidance on the Application of the CLP Criteria**

- Detailed guidance “for experts”;
- On the application of the CLP criteria for physical, health and environmental hazards.

Enable industry to self-classify chemicals and
to provide appropriate hazard communication
information to the target populations.



Guidance on the Application of the CLP Criteria

- Detailed guidance “for experts” -

PART 1: GENERAL PRINCIPLES FOR CLASSIF. AND LABELL.

PART 2: PHYSICAL HAZARDS

PART 3: HEALTH HAZARDS

PART 4: ENVIRONMENTAL HAZARDS

PART 5: ADDITIONAL HAZARDS

ANNEX I: AQUATIC TOXICITY

ANNEX II: RAPID DEGRADATION

ANNEX III: BIOACCUMULATION

ANNEX IV: METALS AND INORGANIC METAL COMPOUNDS

ANNEX V: COLLECTION OF INTERNET LINKS FOR THE USERS OF THE GUIDANCE

ANNEX VI: BACKGROUND TO GUIDANCE FOR SETTING SCLs FOR REPRODUCTIVE TOXICITY

See GHS,
Annex 9, Ch. 7

Exercise

–

substance classification

Classification & Labelling examples

Taken from 4.1.3.4
and 4.1.4.7 of the
EU CLP-Guidance
document

➤ Substances

- A. Hydrophilic substance, straightforward classification based on acute and chronic toxicity data.
- B. Hydrophilic substance, straightforward classification based on acute data, no chronic data available.
- E. “Safety net” classification.

➤ Mixtures

- A & AX. When classification data is available for some or all components of a mixture.
- B2. When information on the classification of the components is available and toxicity data on the mixture as a whole is available for some, but not all three trophic levels.
- C. When no data is available on the mixture or on its components, but test data is available on a similar tested mixture.

Substance example A:

- Hydrophilic substance, straightforward classification based on acute and chronic toxicity data.

For classification assessment, a search of appropriate databases, safety data sheets, C&L-Inventory and other sources of data should be made for the following elements:

ELEMENTS	Value
Physico-chemical properties	
Water solubility / Log Kow:	
Acute aquatic toxicity	
Fish:	
Crustacea	
Algae/aquatic plants	
Chronic aquatic toxicity	
Fish:	
Crustacea:	
Algae/aquatic plants:	
Degradation (evidence of rapid degradation)	
Biotic degradation: Abiotic degradation, hydrolysis: (half-life (d)):	
Bioaccumulation	
Bioconcentration factor (BCF) in fish	

Substance example A, cont.

ELEMENTS	Value
Physico-chemical properties	
Water solubility / Log Kow:	1200 mg/l / 2.75
Acute aquatic toxicity	
Fish:	
<i>Oncorhynchus mykiss</i> :	12 mg/l (96 h LC ₅₀)
<i>Lepomis macrochirus</i> :	2.7 mg/l (96 h LC ₅₀)
Crustacea	
<i>Daphnia magna</i> :	18 mg/l (48 h EC ₅₀)
Algae/aquatic plants	
<i>Scenedesmus subspicatus</i> :	0.056 mg/l (96 h ErC ₅₀)
<i>Lemna gibba</i> :	0.031 mg/l (7 d ErC ₅₀)
Chronic aquatic toxicity	
Fish:	
<i>Danio rerio</i> :	1.2 mg/l (21 d NOEC)
Crustacea:	
<i>Daphnia magna</i> :	1.1 mg/l (21 d NOEC)
Algae/aquatic plants:	
<i>Scenedesmus subspicatus</i> :	0.01 mg/l (96 h NOEC)
Degradation (evidence of rapid degradation)	
Biotic degradation:	86 % in 28 days → RD
Abiotic degradation, hydrolysis: (half-life (d)):	No data
Bioaccumulation	
Bioconcentration factor (BCF) in fish	No data

Acute aquatic hazard

Acute toxicity: ?

≤ 1 mg/l

between 0.01 and 0.1 mg/l

Acute 1, M = 10

Long-term aquatic hazard

Setting M-factors for highly toxic substances (Acute 1 and Chronic 1)

Acute toxicity	M factor
L(E)C ₅₀ value	
$0.1 < L(E)C_{50} \leq 1$	1
$0.01 < L(E)C_{50} \leq 0.1$	10
$0.001 < L(E)C_{50} \leq 0.01$	100
$0.0001 < L(E)C_{50} \leq 0.001$	1000
$0.00001 < L(E)C_{50} \leq 0.0001$	10000
(continue in factor 10 intervals)	

(L(E)C₅₀ and NOEC (or EC_x) in mg/l)

Substance example A, cont.

ELEMENTS	Value
Physico-chemical properties	
Water solubility / Log Kow:	1200 mg/l / 2.75
Acute aquatic toxicity	
Fish:	
<i>Oncorhynchus mykiss</i> :	12 mg/l (96 h LC ₅₀)
<i>Lepomis macrochirus</i> :	2.7 mg/l (96 h LC ₅₀)
Crustacea	
<i>Daphnia magna</i> :	18 mg/l (48 h EC ₅₀)
Algae/aquatic plants	
<i>Scenedesmus subspicatus</i> :	0.056 mg/l (96 h ErC ₅₀)
<i>Lemna gibba</i> :	0.031 mg/l (7 d ErC ₅₀)
Chronic aquatic toxicity	
Fish:	
<i>Danio rerio</i> :	1.2 mg/l (21 d NOEC)
Crustacea:	
<i>Daphnia magna</i> :	1.1 mg/l (21 d NOEC)
Algae/aquatic plants:	
<i>Scenedesmus subspicatus</i> :	0.01 mg/l (96 h NOEC)
Degradation (evidence of rapid degradation)	
Biotic degradation:	86 % in 28 days → RD
Abiotic degradation, hydrolysis: (half-life (d)):	No data
Bioaccumulation	
Bioconcentration factor (BCF) in fish	No data

Acute aquatic hazard

Acute toxicity:

≤ 1 mg/l

between 0.01 and 0.1 mg/l

Acute 1, M = 10

Long-term aquatic hazard

Chronic toxicity: ?

≤ 1 mg/l

between 0.001 and 0.01 mg/l

Degradation:

Rapidly degradable

Chronic

Criteria for Long-term hazard (categories Chronic 1-3)

Adequate chronic toxicity data available	
Non-rapidly degradable (NRD) substance	Rapidly degradable (RD) substances
Category: Chronic 1 NOEC or $EC_x \leq 0.1$	Category: Chronic 1 NOEC or $EC_x \leq 0.01$
Category: Chronic 2 $0.1 < \text{NOEC}$ or $EC_x \leq 1$	Category: Chronic 2 $0.01 < \text{NOEC}$ or $EC_x \leq 0.1$
	Category: Chronic 3 $0.1 < \text{NOEC}$ or $EC_x \leq 1$

➤ Toxicity + degradation

Substance example A, cont.

ELEMENTS	Value
Physico-chemical properties	
Water solubility / Log Kow:	1200 mg/l / 2.75
Acute aquatic toxicity	
Fish:	
<i>Oncorhynchus mykiss</i> :	12 mg/l (96 h LC ₅₀)
<i>Lepomis macrochirus</i> :	2.7 mg/l (96 h LC ₅₀)
Crustacea	
<i>Daphnia magna</i> :	18 mg/l (48 h EC ₅₀)
Algae/aquatic plants	
<i>Scenedesmus subspicatus</i> :	0.056 mg/l (96 h ErC ₅₀)
<i>Lemna gibba</i> :	0.031 mg/l (7 d ErC ₅₀)
Chronic aquatic toxicity	
Fish:	
<i>Danio rerio</i> :	1.2 mg/l (21 d NOEC)
Crustacea:	
<i>Daphnia magna</i> :	1.1 mg/l (21 d NOEC)
Algae/aquatic plants:	
<i>Scenedesmus subspicatus</i> :	0.01 mg/l (96 h NOEC)
Degradation (evidence of rapid degradation)	
Biotic degradation:	86 % in 28 days
Abiotic degradation, hydrolysis: (half-life (d)):	No data
Bioaccumulation	
Bioconcentration factor (BCF) in fish	No data

Acute aquatic hazard

Acute toxicity:

≤ 1 mg/l

between 0.01 and 0.1 mg/l

Acute 1, M = 10

Long-term aquatic hazard

Chronic toxicity:

≤ 1 mg/l

between 0.001 and 0.01 mg/l

Degradation:

Rapidly degradable

Chronic 1, M = 1

Setting M-factors for highly toxic substances


(Acute 1 and Chronic 1)

Acute toxicity	M factor	Chronic toxicity	M factor	
L(E)C ₅₀ value		NOEC value	NRD ^a components	RD ^b components
$0.1 < L(E)C_{50} \leq 1$	1	$0.01 < NOEC \leq 0.1$	1	-
$0.01 < L(E)C_{50} \leq 0.1$	10	$0.001 < NOEC \leq 0.01$	10	1
$0.001 < L(E)C_{50} \leq 0.01$	100	$0.0001 < NOEC \leq 0.001$	100	10
$0.0001 < L(E)C_{50} \leq 0.001$	1000	$0.00001 < NOEC \leq 0.0001$	1000	100
$0.00001 < L(E)C_{50} \leq 0.0001$	10000	$0.000001 < NOEC \leq 0.00001$	10000	1000
(continue in factor 10 intervals)		(continue in factor 10 intervals)		

(L(E)C₅₀ and NOEC (or EC_x) in mg/l)



Labelling elements

- Acute (short-term) aquatic hazard - Categories Acute 1 to 3

	<u>Acute 1</u>	<u>Acute 2</u>	<u>Acute 3</u>
Pictogram		No Pictogram	No Pictogram
Signal word	Warning	No word	No word
Hazard Statement	H400: Very toxic to aquatic life	H401: Toxic to aquatic life	H402: Harmful to aquatic life

Labelling elements

➤ Long-term aquatic hazard - Categories Chronic 1 to 3

	<u>Chronic 1</u>	<u>Chronic 2</u>	<u>Chronic 3</u>
Pictogram			No Pictogram
Signal word	Warning	No word	No word
Hazard Statement	H410: Very toxic to aquatic life with long-lasting effects	H411: Toxic to aquatic life with long-lasting effects	H412: Harmful to aquatic life with long-lasting effects

Safety net Chronic 4 - H413: May cause long lasting harmful effects to aquatic life.

Substance example A, cont.

Aquatic hazard classification

Acute aquatic hazard: **Acute 1, M = 10**

Long-term aquatic hazard: **Chronic 1, M = 1**

Labelling elements based on the classification:

Element	Aquatic hazard information that could appear on the label
GHS Pictogram	GHS09
Signal Word	WARNING
Hazard Statement	H410 (H400, H410 → H410)
Precautionary statement(s)	P273, P391, P501

See hazard communication Table 4.1.4

NOTE! Note that in accordance with GHS hazard statement H400 may be considered redundant and therefore not included on the label because hazard statement H410 also applies. (**GHS 1.4.10.5.3.3 Precedence for allocation of hazard statements**)

Se also: Table 4.1.6 in EU Guidance document

Aquatic hazard classification	Associated hazard statement	Associated hazard statement that could appear on the label
Acute 1	H400	H400
Acute 1 and Chronic 1	H400; H410	H410
Acute 1 and Chronic 2	H400; H411	H410
Acute 1 and Chronic 3	H400; H412	H410
Acute 1 and Chronic 4	H400; H413	H410
Chronic 1	H410	H410
Chronic 2	H411	H411
Chronic 3	H412	H412
Chronic 4	H413	H413

Substance example B:

- Hydrophilic substance, straightforward classification based on acute data, no chronic data available.

Substance example B, cont.

ELEMENTS	Value
Physico-chemical properties	
Water solubility / Log K _{ow} :	1200 mg/l / 2.75
Acute aquatic toxicity	
Fish: <i>Oncorhynchus mykiss</i> : <i>Lepomis macrochirus</i> :	12 mg/l (96 h LC ₅₀) 2.7 mg/l (96 h LC ₅₀)
Crustacea <i>Daphnia magna</i> :	18 mg/l (48 h EC ₅₀)
Algae/aquatic plants <i>Scenedesmus subspicatus</i> : <i>Lemna gibba</i> :	0.056 mg/l (96 h ErC ₅₀) 0.031 mg/l (7 d ErC ₅₀)
Chronic aquatic toxicity	
Fish:	No data
Crustacea:	No data
Algae/aquatic plants:	No data
Degradation (evidence of rapid degradation)	
Biotic degradation: Abiotic degradation, hydrolysis: (half-life (d)):	86 % in 28 days → RD No data
Bioaccumulation	
Bioconcentration factor (BCF) in fish	560

Acute aquatic hazard

Acute toxicity:

≤ 1 mg/l

between 0.01 and 0.1 mg/l

Acute 1, M = 10

Long-term aquatic hazard

Chronic toxicity: ?

-

Criteria for Long-term hazard

Adequate chronic toxicity data available		In absence of adequate chronic toxicity data
Non-rapidly degradable (NRD) substance	Rapidly degradable (RD) substances	ACUTE TOXICITY + NON-RAPIDLY DEGRADABLE and/or BIOACCUMULATIVE
Category: Chronic 1 NOEC or $EC_x \leq 0.1$	Category: Chronic 1 NOEC or $EC_x \leq 0.01$	
Category: Chronic 2 $0.1 < NOEC$ or $EC_x \leq 1$	Category: Chronic 2 $0.01 < NOEC$ or $EC_x \leq 0.1$	
	Category: Chronic 3 $0.1 < NOEC$ or $EC_x \leq 1$	

➤ Toxicity + degradation and/or bioaccumulation

Substance example B, cont.

ELEMENTS	Value
Physico-chemical properties	
Water solubility / Log K _{ow} :	1200 mg/l / 2.75
Acute aquatic toxicity	
Fish: <i>Oncorhynchus mykiss</i> : <i>Lepomis macrochirus</i> :	12 mg/l (96 h LC ₅₀) 2.7 mg/l (96 h LC ₅₀)
Crustacea <i>Daphnia magna</i> :	18 mg/l (48 h EC ₅₀)
Algae/aquatic plants <i>Scenedesmus subspicatus</i> : <i>Lemna gibba</i> :	0.056 mg/l (96 h ErC ₅₀) 0.031 mg/l (7 d ErC ₅₀)
Chronic aquatic toxicity	
Fish:	No data
Crustacea:	No data
Algae/aquatic plants:	No data
Degradation (evidence of rapid degradation)	
Biotic degradation: Abiotic degradation, hydrolysis: (half-life (d)):	86 % in 28 days → RD No data
Bioaccumulation	
Bioconcentration factor (BCF) in fish	560

Acute aquatic hazard

Acute toxicity:

≤ 1 mg/l

between 0.01 and 0.1 mg/l

Acute 1, M = 10

Long-term aquatic hazard

Chronic toxicity:

-

Acute toxicity:

≤ 100 mg/l

≤ 1 mg/l

Degradation:

Rapidly degradable

Bioaccumulation:

BCF > 500 (Log K_{ow} ≤ 4)

Chronic 1

Long-term hazard in absence of adequate chronic toxicity data

**ACUTE TOXICITY TO FISH
CRUSTACEA OR ALGAE**

≤ 1 mg/l

+

**NON-RAPIDLY
DEGRADABLE**



CHRONIC 1

**ACUTE TOXICITY TO FISH
CRUSTACEA OR ALGAE**

> 1 to ≤ 10 mg/l

+

AND/OR



CHRONIC 2

**ACUTE TOXICITY TO FISH
CRUSTACEA OR ALGAE**

> 10 to ≤ 100 mg/l

+

**BIOACCUMULATIVE
(measured
or potential for
bioaccumulation)**



CHRONIC 3

Substance example B, cont.

ELEMENTS	Value
Physico-chemical properties	
Water solubility / Log K _{ow} :	1200 mg/l / 2.75
Acute aquatic toxicity	
Fish:	
<i>Oncorhynchus mykiss</i> :	12 mg/l (96 h LC ₅₀)
<i>Lepomis macrochirus</i> :	2.7 mg/l (96 h LC ₅₀)
Crustacea	
<i>Daphnia magna</i> :	18 mg/l (48 h EC ₅₀)
Algae/aquatic plants	
<i>Scenedesmus subspicatus</i> :	0.056 mg/l (96 h ErC ₅₀)
<i>Lemna gibba</i> :	0.031 mg/l (7 d ErC ₅₀)
Chronic aquatic toxicity	
Fish:	No data
Crustacea:	No data
Algae/aquatic plants:	No data
Degradation (evidence of rapid degradation)	
Biotic degradation:	86 % in 28 days → RD
Abiotic degradation, hydrolysis: (half-life (d)):	No data
Bioaccumulation	
Bioconcentration factor (BCF) in fish	560

Acute aquatic hazard

Acute toxicity:

≤ 1 mg/l

between 0.01 and 0.1 mg/l

Acute 1, M = 10

Long-term aquatic hazard

Chronic toxicity:

-

Acute toxicity:

≤ 100 mg/l

≤ 1 mg/l

between 0.01 and 0.1 mg/l

Degradation:

Rapidly degradable

Bioaccumulation:

BCF > 500 (Log K_{ow} ≤ 4)

Chronic 1, M = 10

Setting M-factors for highly toxic substances

(Acute 1 and Chronic 1)

Acute toxicity	M factor	Chronic toxicity	M factor	
L(E)C ₅₀ value		NOEC value	NRD ^a components	RD ^b components
$0.1 < L(E)C_{50} \leq 1$	1	$0.01 < NOEC \leq 0.1$	1	-
$0.01 < L(E)C_{50} \leq 0.1$	10	$0.001 < NOEC \leq 0.01$	10	1
$0.001 < L(E)C_{50} \leq 0.01$	100	$0.0001 < NOEC \leq 0.001$	100	10
$0.0001 < L(E)C_{50} \leq 0.001$	1000	$0.00001 < NOEC \leq 0.0001$	1000	100
$0.00001 < L(E)C_{50} \leq 0.0001$	10000	$0.000001 < NOEC \leq 0.00001$	10000	1000
(continue in factor 10 intervals)		(continue in factor 10 intervals)		

(L(E)C₅₀ and NOEC (or EC_x) in mg/l)

Substance example B, cont.

Aquatic hazard classification

Acute aquatic hazard: **Acute 1, M = 10**

Long-term aquatic hazard: **Chronic 1, M = 10**

Labelling elements based on the classification:

Element	Aquatic hazard information that could appear on the label
GHS Pictogram	GHS09
Signal Word	WARNING
Hazard Statement	H410 (H400, H410 → H410)
Precautionary statement(s)	P273, P391, P501

Criteria for
environmental hazard
classification
-
mixtures

Substance ingredients

- It is important to get a clear picture on which substances are contained in a mixture.
- Basic information would include: (i) the substance identity, (ii) its classification (iii) any applied M-factor, and (iv) concentration in the mixture.
- Where an **ingredient in a mixture is itself a mixture**, it is generally necessary to get information on the ingredient substances of the first mixture.

NOTE! Further dialogue with the supplier may be necessary to obtain additional information.

Suppliers in a supply chain shall cooperate to meet the requirements for classification, labelling and packaging – CLP, Art. 4.9

Testing of mixtures must be avoided !

- Testing of mixtures is highly complex. Both in conduct of the test, and in interpretation of data.
- Alternative approaches such as the **summation method, should be considered**, particularly where testing would involve the use of vertebrate animals such as fish.

NOTE! *Degradability and bioaccumulation* tests for mixtures are not used as they are usually difficult to interpret, and such tests may be meaningful only for single substances.

Classification of mixtures

➤ The approach used is dependent upon the type of information available for the mixture itself and for its components.

- Criteria **as for substances** – Using data on the mixture itself;
- **Bridging principles** - Data on similar tested mixtures; or
- **The Summation method** – Classification based on individual ingredients.

However: Testing of mixtures must be avoided !

It is **generally the summation** of the quantities of the hazardous components **that should be used** to determine a specific hazard classification of the mixture.

Summation method

➤ Short-term (acute) hazard:

Summation of components:	Mixture is classified as:
$\Sigma(\text{Acute } 1 \times M) \geq 25 \%$	Acute 1

➤ Long-term aquatic hazard (**a stepwise procedure**):

Summation of components:	Mixture is classified as:
$\Sigma(\text{Chronic } 1 \times M) \geq 25 \%$	Chronic 1
$\Sigma(\text{Chronic } 1 \times M \times 10) + \Sigma(\text{Chronic } 2) \geq 25 \%$	Chronic 2
$\Sigma(\text{Chronic } 1 \times M \times 100) + \Sigma(\text{Chronic } 2 \times 10) + \Sigma(\text{Chronic } 3) \geq 25 \%$	Chronic 3
$\Sigma(\text{Chronic } 1) + \Sigma(\text{Chronic } 2) + \Sigma(\text{Chronic } 3) + \Sigma(\text{Chronic } 4) \geq 25 \%$	Chronic 4 (Safety-net)

Exercise

–

mixture classification


(principle use of the
Summation method)

Summation method, example on Long term effects step 1:

Mixture classified as Category Chronic 1 if

$$\sum(\text{Chronic Category 1} \times M) \geq 25\%$$

	Ingr. A	Ingr. B	Ingr. C
	<u>1%</u>	<u>10%</u>	<u>10%</u>
Chronic 3			
Chronic 2			●
Chronic 1, M1		●	
Chronic 1, M10	●		
Chronic 1, M100			




$$10\% + 10\% = 20\%, \text{ which is } < 25\%.$$

Hence, mixture not classified as Chronic 1.

Summation method, example on Long term effects step 2:

Mixture classified as Category Chronic 2 if

$$\sum(\text{Category Chronic 1} \times M \times 10) + \sum(\text{Category Chronic 2}) \geq 25\%$$



100% + 100% + 10% = 210%, which is $\geq 25\%$.
Hence, mixture classified as Chronic 2.

Classification & Labelling examples

Taken from 4.1.3.4
and 4.1.4.7 of the
EU CLP-Guidance
document

➤ Substances

- A. Hydrophilic substance, straightforward classification based on acute and chronic toxicity data.
- B. Hydrophilic substance, straightforward classification based on acute data, no chronic data available.
- E. “Safety net” classification.

➤ Mixtures

- A & AX. When classification data is available for some or all components of a mixture.
- B2. When information on the classification of the components is available and toxicity data on the mixture as a whole is available for some, but not all three trophic levels.
- C. When no data is available on the mixture or on its components, but test data is available on a similar tested mixture.

Mixture example A:

- When classification data is available for some or all components of a mixture.

Mixture example A, cont.

Information on ingredients classification and concentration					
	Acute aquatic hazard	M	Long term aquatic hazard	M	C (%)
Astralamid	Acute 1	10	Chronic 1	10	1
Bastralamid	Acute 1	1	Chronic 2	-	3
Castralamid	Not classified	-	Chronic 2	-	10
Dastralamid	Not classified	-	Chronic 3	-	10
Estralamid	Not classified	-	Not classified	-	10
Festralamid	Not classified	-	Not classified	-	66

M = M-factor; C = Concentration

Aquatic hazard classification

Acute aquatic hazard:

Long-term aquatic hazard:

Mixture example A, cont.

Information on ingredients classification and concentration

	Acute aquatic hazard	M	Long term aquatic hazard	M	C (%)
Astralamid	Acute 1	10	Chronic 1	10	1
Bastralamid	Acute 1	1	Chronic 2	-	3
Castralamid	Not classified	-	Chronic 2	-	10
Dastralamid	Not classified	-	Chronic 3	-	10
Estralamid	Not classified	-	Not classified	-	10
Festralamid	Not classified	-	Not classified	-	66

M = M-factor; C = Concentration

Aquatic hazard classification ?



Acute aquatic hazard: **Not classified.**

Long-term aquatic hazard:

Classify for acute hazard if: $\sum (\text{Acute } 1 \times M) \geq 25\%$

- Using the classification of the components of the mixture:
 $(1 \times 10) + (3 \times 1) = 13$ (which is $< 25\%$).

Mixture example A, cont.

Information on ingredients classification and concentration					
	Acute aquatic hazard	M	Long term aquatic hazard	M	C (%)
Astralamid	Acute 1	10	Chronic 1	10	1
Bastralamid	Acute 1	1	Chronic 2	-	3
Castralamid	Not classified	-	Chronic 2	-	10
Dastralamid	Not classified	-	Chronic 3	-	10
Estralamid	Not classified	-	Not classified	-	10
Festralamid	Not classified	-	Not classified	-	66

M = M-factor; C = Concentration

Aquatic hazard classification ?

Acute aquatic hazard: **Not classified.**



Long-term aquatic hazard: **Category Chronic 2.**

Using the classification of the components of the mixture:

- Step 1: $(1 \times 10) = 10\%$ (which is $< 25\% \rightarrow$ Step 2).
- Step 2: $(10 \times 1 \times 10) + 3 + 10 = 113\%$ (which is $\geq 25\%$).

Hence, **classify as Category Chronic 2.**

Mixture example A, cont.

Aquatic hazard classification

Acute aquatic hazard: **Not classified.**

Long-term aquatic hazard: **Category Chronic 2.**

Labelling elements based on the classification:

Element	Aquatic hazard information that could appear on the label
GHS Pictogram	GHS09
Signal Word	-
Hazard Statement	H411
Precautionary statement(s)	P273, P391, P501

Mixture example AX:

Not in the Guidance Document

- When classification data is available for some or all components of a mixture.

Mixture example AX, cont.

Information on ingredients classification and concentration					
	Acute aquatic hazard	M	Long term aquatic hazard	M	C (%)
Substance 1	Acute 1	100	Chronic 1	10	0.2
Substance 2	Acute 1	1	Chronic 2	-	7
Substance 3	Not classified	-	Chronic 2	-	10
Substance 4	Not classified	-	Not classified	-	82.8

M = M-factor; C = Concentration

Aquatic hazard classification ?

➔ Acute aquatic hazard: **Category Acute 1.**

Long-term aquatic hazard:

Classify for acute hazard if: $\sum (\text{Acute } 1 \times M) \geq 25\%$

- **Using the classification of the components of the mixture:**
 $(0.2 \times 100) + (7 \times 1) = 27\%$ (which is $\geq 25\%$).

Mixture example AX, cont.

Information on ingredients classification and concentration					
	Acute aquatic hazard	M	Long term aquatic hazard	M	C (%)
Substance 1	Acute 1	100	Chronic 1	10	0.2
Substance 2	Acute 1	1	Chronic 2	-	7
Substance 3	Not classified	-	Chronic 2	-	0.7
Substance 4	Not classified	-	Not classified	-	82.8

M = M-factor; C = Concentration

Aquatic hazard classification ?


Acute aquatic hazard: **Category Acute 1.**



Long-term aquatic hazard:

Cut-off values

as in Article 11

- Cut-off values are the **minimum concentrations for a substance to be taken into account** for classification purposes.
- These substances are sometimes referred to as relevant ingredients or relevant components.
- The following cut-off values are associated with each substance classification:
 - Acute 1 → 0.1%/M
 - Chronic 1 → 0.1%/M
 - Chronic 2 → 1% 
 - Chronic 3 → 1%
 - Chronic 4 → 1%


Mixture example AX, cont.

Information on ingredients classification and concentration					
	Acute aquatic hazard	M	Long term aquatic hazard	M	C (%)
Substance 1	Acute 1	100	Chronic 1	10	0.2
Substance 2	Acute 1	1	Chronic 2	-	7
Substance 3	Not classified	-	Chronic 2	-	0.7
Substance 4	Not classified	-	Not classified	-	82.8

M = M-factor; C = Concentration

Aquatic hazard classification

Acute aquatic hazard: **Category Acute 1.**

 Long-term aquatic hazard: **Category Chronic 2.**

Using the classification of the components of the mixture:

Step 1: $(0.2 \times 10) = 2\%$ (which is $< 25\%$ → Step 2).

Step 2: $(10 \times 0.2 \times 10) + 7 = 27\%$ (which is $\geq 25\%$).

Mixture example AX, cont.

Aquatic hazard classification

Acute aquatic hazard: **Category Acute 1.**

Long-term aquatic hazard: **Category Chronic 2.**

See hazard
communication
Table 4.1.4

Labelling elements based on the classification:

Element	Aquatic hazard information that could appear on the label
GHS Pictogram	GHS09
Signal Word	WARNING
Hazard Statement	H410 (H400, H411 → H410)
Precautionary statement(s)	P273, P391, P501

NOTE! If classified within several hazard classes or differentiations for a hazard class, all hazard statements resulting from the classification shall appear on the label, unless there is evident duplication or redundancy. (EU CLP Art. 27)

→ This has been interpreted in the CLP-Guidance document that the hazard statements resulting from acute and long-term aquatic hazard classification can be combined on the label. (See table 4.1.6 of the Guidance document).

➤ **Table 4.1.6 (of the Guidance document)**

Aquatic hazard classification	Associated hazard statement	Associated hazard statement that could appear on the label
Acute Category 1	H400	H400
Acute 1 and Chronic 1	H400; H410	H410
Acute 1 and Chronic 2	H400; H411	H410
Acute 1 and Chronic 3	H400; H412	H410
Acute 1 and Chronic 4	H400; H413	H410
Chronic Category 1	H410	H410
Chronic Category 2	H411	H411
Chronic Category 3	H412	H412
Chronic Category 4	H413	H413

Mixture example B2:

- When information on the classification of the components is available and toxicity data on the mixture as a whole is available for some, but not all three trophic levels.

Mixture example B2, cont.

Information on components classification and concentration

	Acute aquatic hazard	M	Long-term aquatic hazard	M	C (%)
Frustrin	Acute 1	1	Chronic 1	1	40
Gladobrin	Acute 1	1	Chronic 3	-	60

Acute (short-term) aquatic toxicity	Value	Test method ((EC) No. 440/2008) or OECD guideline / remarks
Algae/aquatic plants: Mixture (<i>Scenedesmus subspicatus</i>)	15 mg/l (72 or 96 hr ErC ₅₀)	C.3 / static, GLP
Chronic (long-term) aquatic toxicity		
Algae/aquatic plants: Mixture (<i>Scenedesmus subspicatus</i>)	1.5 mg/l (96 h NOEC)	C.3 / static, GLP

Aquatic hazard classification ?



Acute aquatic hazard: **Category Acute 1.**



Long-term aquatic hazard: **Category Chronic 1.**

Mixture example B2, cont.

Aquatic hazard classification

Acute aquatic hazard: **Category Acute 1.**

Long-term aquatic hazard: **Category Chronic 1.**

See hazard
communication
Table 4.1.4

Labelling elements based on the classification:

Element	Aquatic hazard information that could appear on the label
GHS Pictogram	GHS09
Signal Word	WARNING
Hazard Statement	H410 (H400, H410 → H410)
Precautionary statement(s)	P273, P391, P501

Substance example E:

- “Safety net” classification.

Substance example E, cont.

ELEMENTS	Value
Physico-chemical properties	
Water solubility / Log K _{ow} :	0.009 mg/l / 5.4
Acute aquatic toxicity	
Fish:	No data
Crustacea <i>Daphnia magna</i> :	> 1 mg/l (48 h EC ₅₀) (Static, nominal concentr., non-GLP)
Algae/aquatic plants	No data
Chronic aquatic toxicity	
Fish:	No data
Crustacea:	No data
Algae/aquatic plants:	No data
Degradation (evidence of rapid degradation)	
Biotic degradation:	No data
Abiotic degradation, hydrolysis: (half-life (d)):	No data
Bioaccumulation	
Bioconcentration factor (BCF) in fish	No data

Acute aquatic hazard

Acute toxicity:

No sufficient data

- Not classified due to lack of data

Long-term aquatic hazard

Chronic toxicity:

No data

Degradation:

Not rapidly degradable
(by default in absence of measured data)

Bioaccumulation:

Log K_{ow} > 4

Substance example E, cont.

ELEMENTS	Value
Physico-chemical properties	
Water solubility / Log K _{ow} :	0.009 mg/l / 5.4
Acute aquatic toxicity	
Fish:	No data
Crustacea <i>Daphnia magna</i> :	> 1 mg/l (48 h EC ₅₀) (Static, nominal concentr., non-GLP)
Algae/aquatic plants	No data
Chronic aquatic toxicity	
Fish:	No data
Crustacea:	No data
Algae/aquatic plants:	No data
Degradation (evidence of rapid degradation)	
Biotic degradation:	No data
Abiotic degradation, hydrolysis: (half-life (d)):	No data
Bioaccumulation	
Bioconcentration factor (BCF) in fish	No data

$$S_w \leq 1 \text{ mg/l}$$

Acute aquatic hazard

Acute toxicity:

No sufficient data

- Not classified due to lack of data

Long-term aquatic hazard

Chronic toxicity:

No data

Degradation:

Not rapidly degradable
(by default in absence of measured data)

Bioaccumulation:

Log K_{ow} > 4

Chronic 4
(Safety net)

Substance example E, cont.

Aquatic hazard classification

Acute aquatic hazard: **Not classified due to lack of data**

Long-term aquatic hazard: **Chronic 4**

Labelling elements based on the classification:

Element	Aquatic hazard information that could appear on the label
GHS Pictogram	-
Signal Word	-
Hazard Statement	H413
Precautionary statement(s)	P273, P501

***Thank You
for Your Attention***